Non-principal orders in algebraic number fields with half-factorial localizations

Andreas Philipp

Institute for Mathematics and Scientific Computing
Karl-Franzens University Graz

October 16, 2009 – Integers Conference
University of West Georgia
Carrollton, Georgia

This work is supported by the FWF Austrian Science Found (FWF Grant P18779-N13).

Table of contents

1. Introduction
   - Example of non-unique factorization
   - Basic facts about non-unique factorization in orders
   - Some definitions
   - Important invariants
2. Connection between rings and monoids
   - Translation of the problem to a combinatorial problem
   - Block monoids
3. Maximal orders
4. Non-principal orders
5. Main result
   - Easy consequences

Example of non-unique factorization

Factorization in rings of algebraic integers is not necessarily unique.
Example: $\mathbb{Z}[3i] \subset \mathbb{Q}(i)$
The elements $3+6i$, $3-6i$, 3, 5 are all irreducible.

$$(3+6i)(3-6i) = 3^2 \cdot 5$$

This phenomenon is called non-unique factorization.

The phenomenon of non-unique factorization is purely multiplicative.

Non-unique factorization in maximal orders

Let K be an algebraic number field, $\mathcal{O}_K$ be the ring of integers, i.e. the maximal order, of K.
Despite for the integers $\mathbb{Z}$ factorization in such a ring $\mathcal{O}_K$ needs no longer to be unique.
But these rings $\mathcal{O}_K$ are integrally closed and indeed they are Dedekind domains. Thus their (arithmetic) structure can be described well in the terms of classical ideal theory.

Non-unique factorization in non-principal orders

Non-principal orders (subrings $\mathcal{O} \subset \mathcal{O}_K$) are not integrally closed anymore. In particular, $\mathcal{O}$ is never factorial.

The arithmetic of a non-principal order depends on the Picard group, on the localizations at singular primes and on a yet not understood interplay between these data.

Introduction

Let K be an algebraic number field, $\mathcal{O}_K$ be the maximal order in K, and let $\mathcal{P} = \{p_1, \ldots, p_s\}$ be a set of primes that are inert in $\mathcal{O}_K$ (that is, $p_1, \ldots, p_s$ are prime in $\mathcal{O}_K$).
Then

$$\mathcal{O} = \mathbb{Z} + p_1 \cdot \ldots \cdot p_s \mathcal{O}_K$$

is a non-principal order in K such that all localizations are half-factorial.

What about the arithmetic of such a non-principal order?

$$\left.\begin{array}{ll} \triangle(\mathcal{O}) & \min \triangle(\mathcal{O}) \\ \rho(\mathcal{O}) & \mathsf{c}(\mathcal{O}) \end{array}\right\} \quad \text{invariants characterizing the arithmetic}$$

These invariants measure the deviation from unique-factorization.

$$\text{unique-factorization} \;\Leftrightarrow\; \triangle(\mathcal{O}) = \emptyset,\ \min \triangle(\mathcal{O}) = 0,\ \rho(\mathcal{O}) = 1,\ \mathsf{c}(\mathcal{O}) = 0$$

Some definitions

By a *monoid* we always mean a commutative semigroup with identity which satisfies the cancellation law (that is, if a, b, $c \in H$ with $ab = ac$, then $b = c$ follows).

Let H be a monoid. We denote by $H^\times$ the set of invertible elements of H, and we say that H is *reduced* if $H^\times = \{1\}$.
Let $H_{\text{red}} = H/H^\times = \{aH^\times | a \in H\}$ be the associated reduced monoid, and $\mathsf{q}(H)$ a (the) quotient group of H.

Let R be a domain. Then $(R^\bullet = R\backslash\{0\}, \cdot)$ is a monoid, and $(R^\bullet)_{\text{red}}$ is isomorphic to the monoid of nonzero principal ideals $\mathcal{H}(R) = \{aR | a \in R^\bullet\}$.

Factorizations

Let H be an atomic monoid and $a \in H \setminus H^\times$.
A *factorization* of a (in H) is a decomposition of a into a product of irreducible elements (atoms), that is

$$a = u_1 \cdot \ldots \cdot u_n \quad \text{for } n \in \mathbb{N} \text{ and } u_1, \ldots, u_n \in \mathcal{A}(H)$$

Then this n is called a *length* of a (in H).

The set

$$\mathsf{L}(a) = \{n \in \mathbb{N} | n \text{ is a length of } a\}$$

is called the *set of lengths (of a)*.

We call H *half-factorial* if $|\mathsf{L}(a)| = 1$ for all $a \in H \setminus H^\times$.

Set of distances

For a finite subset $L = \{a_1, \ldots, a_t\} \subset \mathbb{Z}$ $(a_1 < a_2 < \ldots < a_t)$ let

$$\triangle(L) = \{a_{\nu+1} - a_\nu | \nu \in [1, t-1]\} \subset \mathbb{N}$$

denote the *set of (successive) distances* of L. Then

$$\triangle(H) = \bigcup_{a \in H} \triangle(\mathsf{L}(a)) \subset \mathbb{N}$$

denotes the *set of distances of* H.
Clearly, H is half-factorial if and only if $\triangle(H) = \emptyset$.

We call $\min \triangle(H)$ the *minimum distance* of H and we set $\min \triangle(H) = 0$ if $\triangle(H) = \emptyset$.

Translation of the problem to a combinatorial problem

Let $\mathcal{O}$ be a non-principal order in an algebraic number field K, $\mathcal{O}_K$ be the corresponding maximal order, $\mathfrak{f} = (\mathcal{O} : \mathcal{O}_K)$ be the conductor, $\mathcal{P} = \{\mathfrak{p} \in \mathfrak{X}(\mathcal{O}) | \mathfrak{p} \not\supset \mathfrak{f}\}$, $\mathcal{P}^* = \{\mathfrak{p} \in \mathfrak{X}(\mathcal{O}) | \mathfrak{p} \supset \mathfrak{f}\}$ and $T = \prod_{\mathfrak{p} \in \mathcal{P}^*} (\mathcal{O}_\mathfrak{p}^\bullet)_{\mathrm{red}}$.
We have the following isomorphisms

$$\mathcal{I}^*(\mathcal{O}) \tilde{\to} \coprod_{\mathfrak{p} \in \mathfrak{X}(\mathcal{O})} (\mathcal{O}_\mathfrak{p}^\bullet)_{\mathrm{red}} \tilde{\to} \mathcal{F}(\mathcal{P}) \times T$$

The diagonal embedding induces a cofinal divisor homomorphism

$$\varphi : \mathcal{O}^\bullet \to \coprod_{\mathfrak{p} \in \mathfrak{X}(\mathcal{O})} (\mathcal{O}_\mathfrak{p}^\bullet)_{\mathrm{red}} \tilde{\to} \mathcal{F}(\mathcal{P}) \times T$$

and we set $H = \varphi(\mathcal{O}^\bullet)$.
Then $H \cong (\mathcal{O}^\bullet)_{\mathrm{red}}$ and $H \subset \mathcal{F}(\mathcal{P}) \times T$ is a saturated and cofinal submonoid and $\mathrm{Pic}(\mathcal{O}) = \mathcal{C}(\varphi) = (\mathcal{F}(\mathcal{P}) \times T)/H$.
We set $\mathrm{Pic}(\mathcal{O}) = G$. We identify these groups.

Definition

Let G be an additive abelian group, $\mathcal{F}(G)$ be the free abelian monoid with bases G, and $\sigma : \mathcal{F}(G) \to G$ the unique homomorphism satisfying $\sigma(g) = g$ for all $g \in G$.
We call $\mathcal{B}(G) = \{S \in \mathcal{F}(G) | \sigma(S) = 0\}$ the block monoid over G.
This is the monoid of zero-sum sequences over G.

Block monoids

$$\mathcal{B}(G) \subset \mathcal{B}(H) \subset \mathcal{F}(G) \times T$$

The canonical map

$$\beta_{\mathcal{O}} : \begin{cases} \mathcal{O}^\bullet & \to & \mathcal{B}(H) \\ a & \mapsto & \left(\prod_{\mathfrak{p}\in\mathcal{P}} [\mathfrak{p}]^{\mathsf{v}_\mathfrak{p}(a)}\right) (a\mathcal{O}_\mathfrak{p}^\times)_{\mathfrak{p}\in\mathcal{P}^*} \end{cases}$$

is a transfer homomorphism.
The arithmetical structures of $\mathcal{O}^\bullet$ and $\mathcal{B}(H)$ are almost identical.

Maximal orders

If $\mathcal{O} = \mathcal{O}_K$, i.e. if $\mathcal{O}$ is a maximal order, then $T = \{1\}$ and $\mathcal{B}(G) = \mathcal{B}(H)$.
In this case the situation is easier.

$$\mathcal{B}(G) \subset \mathcal{F}(G)$$

The canonical map

$$\beta_{\mathcal{O}} : \left\{ \begin{array}{ccc} \mathcal{O}^\bullet & \to & \mathcal{B}(G) \\ a & \mapsto & \left(\prod_{\mathfrak{p} \in \mathcal{P}} [\mathfrak{p}]^{\mathsf{v}_\mathfrak{p}(a)} \right) \end{array} \right.$$

is a transfer homomorphism.
Thus the arithmetical structures of $\mathcal{O}^\bullet$ and $\mathcal{B}(G)$ are almost identical.

$\mathcal{B}(G)$ and $|G|$

Let G be an abelian group and $\mathcal{B}(G)$ be its block monoid.

- $|G| \leq 2$
  $\mathcal{B}(G)$ is half-factorial.
  $\Rightarrow \ \triangle(\mathcal{B}(G)) = \emptyset$
- $|G| \geq 3$
  $1 \in \triangle(\mathcal{B}(G))$
  $\Rightarrow \ \min \triangle(\mathcal{B}(G)) = 1$

Example for $\min \triangle(\mathcal{B}(G))$

Let $|G| \geq 3$ and $\exp(G) = n = 2k + 1 \geq 3$.
Then

$$\left[(2g)g\right]^n = \left[(2g)^n\right]\left[g^n\right] = \left[(2g)^k g\right]^2 \left[(2g)g^{2k-1}\right]$$

Thus $1 \in \triangle(\mathcal{B}(G))$ and therefore $\min \triangle(\mathcal{B}(G)) = 1$.

Maximal orders

Theorem

Let $\mathcal{O}_K$ be the maximal order of an algebraic number field K.
Then

- *$\mathcal{O}_K$ is factorial if and only if $|\operatorname{Pic}(\mathcal{O}_K)| = 1$.*
- *$\mathcal{O}_K$ is half-factorial if and only if $|\operatorname{Pic}(\mathcal{O}_K)| \leq 2$.*

Proof.

The first part was already known by Kummer in the 19th century and the second part by Carlitz in 1960. □

Corollary

Let $\mathcal{O}_K$ be the maximal order of an algebraic number field K.
Then

$$\min \triangle(\mathcal{O}_K) \leq 1$$

But: This does not carry over to non-principal orders.

$\mathcal{B}(G) \subset \mathcal{B}(H)$

$\mathcal{B}(G) \subset \mathcal{B}(H)$ is a divisor closed submonoid.

Immediate consequences

- $\mathcal{A}(\mathcal{B}(G)) \subset \mathcal{A}(\mathcal{B}(H))$, i.e. each atom of $\mathcal{B}(G)$ is an atom of $\mathcal{B}(H)$.
- $\triangle(\mathcal{B}(G)) \subset \triangle(\mathcal{B}(H))$.
- If $|G| \geq 3$ then $1 \in \triangle(\mathcal{B}(G))$ and thus
  - $\min \triangle(\mathcal{B}(H)) = 1$, $\mathsf{c}(H) \geq 3$, and $\rho(H) > 1$

Theorem

Let $\mathcal{O}$ be an order in an algebraic number field and $|\operatorname{Pic} \mathcal{O}| \geq 3$. Then we have

- $\min \triangle(\mathcal{O}) = 1$
- $\mathsf{c}(\mathcal{O}) \geq 3$
- $\rho(\mathcal{O}) > 1$, *i.e. $\mathcal{O}$ is not half-factorial.*

But: What can we say about these invariants if $|\operatorname{Pic}(\mathcal{O})| \leq 2$?

Strategy

$$H \subset \mathcal{F}(\mathcal{P}) \times T \quad \text{saturated}$$

For an order $\mathcal{O}$ with half-factorial localizations let $\mathcal{P}^* = \{\mathfrak{p}_1, \ldots, \mathfrak{p}_r\}$.

$$D_i \cong (\mathcal{O}^\bullet_{\mathfrak{p}_i})_{\mathrm{red}} \quad \text{for all } i = 1, \ldots, r$$

$\Rightarrow$ D_i are half-factorial finitely primary monoids, i.e. they have a nice structure.

Strategy: Use their structure and $G = \mathsf{q}(D/H)$ (=class group) to determine the arithmetic of H.

Theorem

Let $\mathcal{O}$ be an order in an algebraic number field such that all localizations are half-factorial and $|\operatorname{Pic}(\mathcal{O})| \le 2$.
Then we have

1. $\rho(\mathcal{O}) \in \{1, \frac{3}{2}, 2\}$.
2. $\mathsf{c}(\mathcal{O}) \in \{2, 3\}$ *if* $\rho(\mathcal{O}) = 1$.
3. $\mathsf{c}(\mathcal{O}) \ge 3$ *if* $\rho(\mathcal{O}) \ge \frac{3}{2}$.
4. $\mathsf{c}(\mathcal{O}) \ge 4$ *if* $\rho(\mathcal{O}) = 2$.
5. $\min \triangle(\mathcal{O}) \le 1$.

Corollary

Let $\mathcal{O}$ be an order in an algebraic number field such that all localizations are half-factorial.
If all localizations of $\mathcal{O}$ are finitely primary monoids of exponent 1 then the following are equivalent:

1. *$\mathcal{O}$ is half-factorial.*
2. $\mathsf{c}(\mathcal{O}) = 2$.

In particular, for quadratic or cubic number fields the condition is always fulfilled.

Corollary

Let $\mathcal{O}$ be an order in an algebraic number field such that all localizations are half-factorial.
Then

$$\min \triangle(\mathcal{O}) \le 1$$